

# Incubation Period for Neuroinvasive Toscana Virus Infections

Lison Laroche,<sup>1</sup> Frédéric Jourdain,<sup>1</sup> Nazli Ayhan, Anne-Laure Bañuls, Rémi Charrel, Jorian Prudhomme

Toscana virus (TOSV) is an emerging pathogen in the Mediterranean area and is neuroinvasive in its most severe form. Basic knowledge on TOSV biology is limited. We conducted a systematic review on travel-related infections to estimate the TOSV incubation period. We estimated the incubation period at 12.1 days.

**T**oscana virus (TOSV) is an arthropodborne virus transmitted to humans through a bite from an infected sand fly (1). An RNA virus, it belongs to the genus *Phlebovirus*, species (*Sandfly fever Naples phlebovirus* family *Phenuiviridae*, order *Bunyavirales*) (2). TOSV infections are endemic to the Mediterranean basin and are considered frequent even though they are neglected (3). TOSV can be neuroinvasive and is a major cause of meningitis and encephalitis during summer months in areas to which it is endemic (4). However, most infections are asymptomatic or produce mild symptoms (5). Thus, TOSV cases are massively underestimated and unreported. Cases are mainly diagnosed by reverse transcription PCR in cerebrospinal fluid, blood, and, rarely, urine or by detecting virus-specific IgM or IgG (6). A total of 3 different TOSV lineages (A, B, and C) have been identified, but no clear evidence of a link between clinical manifestation and lineages exists (7).

In this study, we considered the incubation period (IP) of an infectious disease as the delay between infection and symptom onset; this definition differs from the latent period, which is defined as the time from infection to infectiousness. For arthropodborne viruses, the infectious bite represents the date of in-

fection (8). The potential period of exposure is represented by the length of stay in the country of infection before symptom onset. We therefore focused on imported cases.

Determining the IP is primordial for disease surveillance, outbreak investigation, public health interventions, infectious disease control, and modeling (9). However, IP estimates are often unsourced, imprecise, and based on limited evidence, as illustrated by the heterogeneous values proposed (Appendix Table 1, <https://wwwnc.cdc.gov/EID/article/27/12/20-3172-App1.pdf>). In this context, we conducted a systematic review of symptomatic travel-related neuroinvasive forms of TOSV to provide an evidence-based estimate of the IP.

## The Study

We used PubMed and ISI Web of Knowledge search engines with no restriction on language and the phrase “Toscana AND virus AND (case report OR case-report OR travel\* OR import\*).” We conducted a systematic search on ProMed and Google Scholar, as well as cross-reference checking. The inclusion criteria were laboratory-documented acute TOSV infection, indication of a travel-related infection in a TOSV-endemic area, and number of days between travel return and symptom onset. Two reviewers screened titles, abstracts, and full-text articles independently.

We extracted clinic and biologic elements from neuroinvasive TOSV case reports. For each patient, data related to the duration of travel and the time of symptom onset, gender, age, country in which case was reported, and country of infection were reported.

To estimate the IP, we used censored time-to-event models (10). Interval-censored observations related to travel duration represented the exposure time. Absence of a departure date was treated as left-censored data, whereas onset of illness during the travel period was considered right-censored. We performed data analysis by using R with the *icenReg*

Author affiliations: Université de Montpellier, Institut de Recherche pour le Développement, Centre National de la Recherche Scientifique, Montpellier, France (L. Laroche, F. Jourdain, A.-L. Bañuls, J. Prudhomme); Santé Publique France, Saint-Maurice, France (F. Jourdain); Aix-Marseille Université, Institut de Recherche pour le Développement, Institut national de la santé et de la recherche médicale, Marseille, France (N. Ayhan, R. Charrel)

DOI: <https://doi.org/10.3201/eid2712.203172>

<sup>1</sup>These authors contributed equally to this article.

package (R Foundation for Statistical Computing, <https://www.r-project.org>). We defined the data distribution with 4 parametric models (log-normal, log-logistic, Gamma, and Weibull). To determine the best model for our distribution, we calculated the Akaike information criterion. We used the nonparametric log-rank test from the interval package to assess the effect of age and gender as covariates. To check the result stability, we performed an additional Bayesian approach, fitting with the Weibull distribution.

Regarding imported case reports, 142 documents were identified on PubMed and 133 on Web of Knowledge. We removed 79 duplicates and excluded 118 records after screening titles and abstracts. A total of 42 articles were eligible after full-text reading. We then selected 22 documents for data extraction. A total of 24 cases were selected (Appendix Table 2, Figure). All travel-associated cases fulfilling the inclusion criteria were neuroinvasive; these cases were diagnosed in a non-TOSV-endemic area after a stay in a proven TOSV-endemic area (Figure 1).


We selected Weibull distribution because it presented the lower Akaike information criterion (Figure 2; Appendix Table 3). The median IP for neuroinvasive forms is estimated to be 12.1 (95% CI 10.2–14.4) days. In 5% of neuroinvasive cases, symptoms will develop by 6.8 (95% CI 3.8–9.9) days after

an infectious bite; symptoms will develop in 95% of cases by 16.8 (95% CI 13.9–21.7) days after the infectious bite. We found no evidence of age or gender effect on the length of the IP ( $p$  value  $>0.05$  by log-rank test). By using Bayesian analysis, we found an IP of 12.1 (95% CI 9.9–14.4) days (data not shown; results same as Figure 2).


## Conclusions

In the literature, the IP values of TOSV are often heterogeneous, unsourced, or without evidence and therefore do not constitute a valid estimate for clinical or infection control decisions. Our literature review identified 24 neurologic cases of TOSV infection. Some travel durations were reported approximately in case reports and were not included in the analysis. All the data used were based on severe neurologic forms of the disease, which required hospitalization soon after the exposure period.

We estimated the median IP of TOSV at 12 (95% CI 10.2–14.4) days. Considering the delay from infection to symptom onset, this value is greater than that for most other arboviruses (11). Our estimate of the IP is evidence-based and relies on data from well-characterized cases. However, cases that cause mildersymptoms, as opposed to neuroinvasive forms of the disease, might have a shorter IP (similar to



**Figure 1.** Geographic distribution of imported neuroinvasive cases of TOSV (n = 24) and countries of origin of infection. TOSV, Toscana virus.



**Figure 2.** Cumulative percentage of Toscana virus cases manifesting with neurologic symptoms by a given day under the estimates for the Weibull parametric distribution ( $n = 24$ ). Red dashed line represents the median estimation of the incubation period. Solid red horizontal line represents the 95% CI of the median. Gray shading indicates the 95% CI of the values.

other arboviruses). Other symptoms associated with paucisymptomatic forms of TOSV might not have been described yet and should be further investigated to improve case definition and diagnosis.

We also cannot exclude infections by other sandfly fever Naples phleboviruses because of cross-reaction risk in serologic analyses due to their close genetic relationships (12). However, the incidence in the population of other genetically similar phleboviruses is lower than TOSV, and TOSV remains the most common cause of neuroinvasive symptoms (3). Knowledge of TOSV genotypes and their aptitude to cause different clinical forms is limited (12). Analyzing this hypothesis was not possible because of the limited amount of available data. In addition to the genotype, other parameters may influence the IP, such as viral strain, patient's immune status, or viremia (9). The amount of virus transmitted during bites (viral load) could also influence the IP and should be further investigated.

In addition, all other cases were diagnosed in countries or regions to which TOSV is not endemic (United States, United Kingdom, Sweden, Germany, Switzerland, Australia, and France). These imported cases represent a risk for emergence in these areas when vectors are established (13), as has been observed for other vectorborne diseases (14). Moreover, sand flies are known to spread in countries or regions to which TOSV is not endemic (15).

Currently, information on TOSV infections is lacking (12). Precise definitions of the IP should

provide more information on the disease epidemiology and on its development in the human host. Moreover, because the IP is a key parameter for disease modeling (9), it would improve our understanding of the disease transmission dynamics. More reports of travel-related cases and standardization of data collection with reliable information (e.g., location and duration of the trips and precise dates of symptom onset) are clearly needed. The IP estimation will be improved with addition of new data.

#### Acknowledgments

We thank Julian Wei-Tze Tang, James Farrugia, Jonas Schmidt-Chanasit, Josef Georg Heckmann, and David Breen for answering our questions by email to clarify information regarding the TOSV case reports. We also thank Eric Elguero for his support in statistical analysis, Heïdi Lançon for the English revision of the manuscript, and Jérôme Depaquit for reviewing the manuscript.

This research was funded by the IRD (Institut de Recherche pour le Développement), CNRS (Centre National de la Recherche Scientifique), UM (Université de Montpellier) and by the by the European Virus Archive Global (EVA-GLOBAL) project that has received funding from the European Union's Horizon 2020-INFRAIA-2019 research and innovation program under grant agreement No 871029. F.J. is financially supported by the SPF (Santé Publique France) through a doctoral fellowship, J.P. by the INFRAVEC2 project (<https://infravec2.eu>), L.L. by a UM doctoral fellowship, and N.A. by a postdoctoral fellowship from the AMU (Aix Marseille University).

## About the Author

Ms. Laroche is a doctoral student at the French National Research Institute for Sustainable Development and the University of Montpellier, France. Her primary research interests are medical entomology, especially the interactions between pathogens (*Phlebovirus*, *Leishmania*), and their vector (the sand fly).

## References

1. Depaquit J, Grandadam M, Fouque F, Andry PE, Peyrefitte C. Arthropod-borne viruses transmitted by Phlebotomine sandflies in Europe: a review. *Euro Surveill*. 2010;15:19507. <https://doi.org/10.2807/ese.15.10.19507-en>
2. Ayhan N, Prudhomme J, Laroche L, Bañuls AL, Charrel RN. Broader geographical distribution of Toscana virus in the Mediterranean region suggests the existence of larger varieties of sand fly vectors. *Microorganisms*. 2020;8:e114. <https://doi.org/10.3390/microorganisms8010114>
3. Charrel RN, Bichaud L, de Lamballerie X. Emergence of Toscana virus in the Mediterranean area. *World J Virol*. 2012;1:135–41. <https://doi.org/10.5501/wjv.v1.i5.135>
4. Charrel RN, Gallian P, Navarro-Marí JM, Nicoletti L, Papa A, Sánchez-Seco MP, et al. Emergence of Toscana virus in Europe. *Emerg Infect Dis*. 2005;11:1657–63. <https://doi.org/10.3201/eid1111.050869>
5. Braitto A, Corbisiero R, Corradini S, Marchi B, Sancasciani N, Fiorentini C, et al. Evidence of Toscana virus infections without central nervous system involvement: a serological study. *Eur J Epidemiol*. 1997;13:761–4. <https://doi.org/10.1023/A:1007422103992>
6. Ergunay K, Kaplan B, Okar S, Akkutay-Yoldar Z, Kurne A, Arsava EM, et al. Urinary detection of Toscana virus nucleic acids in neuroinvasive infections. *J Clin Virol*. 2015;70:89–92. <https://doi.org/10.1016/j.jcv.2015.07.297>
7. Ayhan N, Charrel RN. Of phlebotomines (sandflies) and viruses: a comprehensive perspective on a complex situation. *Curr Opin Insect Sci*. 2017;22:117–24. <https://doi.org/10.1016/j.cois.2017.05.019>
8. Pinggen M, Bryden SR, Pondeville E, Schnettler E, Kohl A, Merits A, et al. Host inflammatory response to mosquito bites enhances the severity of arbovirus infection. *Immunity*. 2016;44:1455–69. <https://doi.org/10.1016/j.immuni.2016.06.002>
9. Chan M, Johansson MA. The incubation periods of Dengue viruses. *PLoS One*. 2012;7:e50972. <https://doi.org/10.1371/journal.pone.0050972>
10. Cowling BJ, Muller MP, Wong IOL, Ho LM, Louie M, McGeer A, et al. Alternative methods of estimating an incubation distribution: examples from severe acute respiratory syndrome. *Epidemiology*. 2007;18:253–9. <https://doi.org/10.1097/01.ede.0000254660.07942.fb>
11. Rudolph KE, Lessler J, Moloney RM, Kmush B, Cummings DAT. Incubation periods of mosquito-borne viral infections: a systematic review. *Am J Trop Med Hyg*. 2014;90:882–91. <https://doi.org/10.4269/ajtmh.13-0403>
12. Maes P, Adkins S, Alkhovsky SV, Avšič-Zupanc T, Ballinger MJ, Bente DA, et al. Taxonomy of the order Bunyavirales: second update 2018. *Arch Virol*. 2019;164:927–41. <https://doi.org/10.1007/s00705-018-04127-3>
13. Knechtli R, Jenni L. Distribution and relative density of three sandfly (Diptera: Phlebotominae) species in southern Switzerland [in French]. *Ann Parasitol Hum Comp*. 1989;64:53–63. <https://doi.org/10.1051/parasite/198964153>
14. Liang G, Gao X, Gould EA. Factors responsible for the emergence of arboviruses; strategies, challenges and limitations for their control. *Emerg Microbes Infect*. 2015;4:e18. <https://doi.org/10.1038/emi.2015.18>
15. Maroli M, Feliciangeli MD, Bichaud L, Charrel RN, Gradoni L. Phlebotomine sandflies and the spreading of leishmaniases and other diseases of public health concern. *Med Vet Entomol*. 2013;27:123–47. <https://doi.org/10.1111/j.1365-2915.2012.01034.x>

Address for correspondence: Lison Laroche, UMR MIVEGEC, IRD–CNRS–Université de Montpellier, 911 avenue Agropolis, F34394 Montpellier, France; email: lisonlaroche14@gmail.com

# Incubation Period for Neuroinvasive Toscana Virus Infections

## Appendix

**Appendix Table 1.** Estimation of Toscana virus incubation period described in the published literature

Estimation of incubation period	Year	Reference
≥5 days	1996	(1)
From a few days to ≤15 days	1998	(2)
From a few days to 2 weeks	2005	(3)
3–7 days	2005	(4)
3–6 days	2009	(5)
A short incubation period	2013	(6)
Ranges from 3–6 days to 2 weeks	2011	(7)
A short incubation period	2013	(8)
3–7 days	2013	(9)
A short incubation period	2014	(10)
A short incubation period	2014	(11)
A short incubation period (2–7 days)	2014	(12)
From a few days to 2 weeks	2014	(13)
3–7 days	2015	(14)
3–7 days	2015	(15)
From a few days to 2 weeks	2016	(16)
Variable, ranging from a few days to 2–3 weeks	2016	(17)
3–6 days	2016	(18)
≤2 weeks	2017	(19)
3–7 days with a maximum of 2 weeks	2017	(20)
From a few days to approximately 2 weeks	2019	(21)
Usually 3–7 days	2020	(22)

**Appendix Table 2.** Toscana virus case reports (n = 22) included in the analysis of incubation period estimation\*

Reference	Case	Sex	Age, y	Country of infection	Reporting country	Diagnostic methods			Length of stay, d	Time between return and symptom onset, d
						Molecular identification	IgM	IgG		
(23)		M	51	Italy	Australia	+	None	None	17	0
(24)		M	66	Italy	United States	–	+	+	16	5
(25)		M	61	Italy	Switzerland	+	+	+	21	5
(26)†		F	49	Italy	France (Paris)‡	None§	+	+	12	5
(27)	1	M	19	France	Germany	None	+	+	14	12
	2	F	73	Greece	Germany	None	+	+	20	12
(28)		M	43	Italy	Switzerland	None	+	+	NA	7
(29)		M	40	Portugal	Sweden	None	+	+	14	5
(30)†		M	68	Italy	France (Paris)‡	None	+	+	11	9
(31)		M	17	Italy	Switzerland	+	+	+	21	14
(32)		M	82	Italy	United States	None	+	+	14	2
(33)		F	69	Italy	Germany	None	+	+	14	–3
(34)		M	51	Italy	United States	–	+	+	NA	10
(35)		F	18	Italy	United Kingdom	+	–	–	NA	7
(36)		M	65	Italy	United States	+	None	None	21	2
(37)		F	80	Spain	Germany	–	+	+	NA	4
(38)		M	21	Italy	Germany	–	+	+	14	6
(39)		M	65	Italy	United Kingdom	+	None	+	4	12
(40)		M	34	Turkey	Germany	None	+	+	NA	5
(41)	1	M	40	Malta	Switzerland	None	+	+	14	2
	2	F	50	Malta	Switzerland	None	+	+	14	2
(42)		M	53	Italy	Germany	None	+	+	10	–4
(43)		M	20	Italy	Switzerland	–	+	+	13	1
(44)		M	23	Italy	Switzerland	+	+	+	4	10

\*NA, no data available; –, negative to Toscana virus; +, positive to Toscana virus.

†Toscana virus case reports diagnosed in Toscana virus–endemic country but in a non-Toscana virus–endemic region.

‡Proven Toscana virus–endemic country but the case was diagnosed in a non-Toscana virus–endemic area of the country.


§Indicates method was not mentioned in the case report.



**Appendix Table 3.** Incubation estimates, in days, for Toscana virus for 4 competing models (Weibull, Gamma, log-logistic and log-normal) sorted according to the Akaike information criterion\*

Model	Median (95% CI)	Log-likelihood	AIC
Weibull	12.1 (10.2–14.4)	–10.16907	24.33814
Gamma	11.5 (9.3–13.9)	–11.26049	26.52098
Log-logistic	11.9 (9.8–14.4)	–11.50064	27.00128
Log-normal	11.3 (9.3–13.9)	–11.78880	27.57760

\*AIC, Akaike information criterion.



**Appendix Figure.** PRISMA flow diagram adapted from Moher et al. (45) in study of incubation period of Toscana virus

## References

1. Nicoletti L, Ciufolini MG, Verani P. Sandfly fever viruses in Italy. In: Schwarz TF, Siegl G, editors. Imported virus infections. Vienna (Austria): Springer; 1996. p. 41–47.

2. Braito A, Corbisiero R, Corradini S, Fiorentini C, Ciufolini MG. Toscana virus infections of the central nervous system in children: a report of 14 cases. *J Pediatr.* 1998;132:144–8. [PubMed](#)  
[https://doi.org/10.1016/S0022-3476\(98\)70500-1](https://doi.org/10.1016/S0022-3476(98)70500-1)
3. Charrel RN, Gallian P, Navarro-Marí JM, Nicoletti L, Papa A, Sánchez-Seco MP, et al. Emergence of Toscana virus in Europe. *Emerg Infect Dis.* 2005;11:1657–63. [PubMed](#)  
<https://doi.org/10.3201/eid1111.050869>
4. Defuentes G, Rapp C, Imbert P, Durand J-P, Debord T. Acute meningitis owing to phlebotomus fever Toscana virus imported to France. *J Travel Med.* 2005;12:295–6. [PubMed](#)  
<https://doi.org/10.2310/7060.2005.12512>
5. Sonderegger B, Hächler H, Dobler G, Frei M. Imported aseptic meningitis due to Toscana virus acquired on the island of Elba, Italy, August 2008. *Euro Surveill.* 2009;14:19079. [PubMed](#)  
<https://doi.org/10.2807/es.14.01.19079-en>
6. Nougairede A, Bichaud L, Thiberville SD, Ninove L, Zandotti C, de Lamballerie X, et al. Isolation of Toscana virus from the cerebrospinal fluid of a man with meningitis in Marseille, France, 2010. *Vector Borne Zoonotic Dis.* 2013;13:685–8. [PubMed](#) <https://doi.org/10.1089/vbz.2013.1316>
7. Serata D, Rapinesi C, Del Casale A, Simonetti A, Mazzarini L, Ambrosi E, et al. Personality changes after Toscana virus (TOSV) encephalitis in a 49-year-old man: A case report. *Int J Neurosci.* 2011;121:165–9. [PubMed](#) <https://doi.org/10.3109/00207454.2010.537412>
8. Gonen OM, Sacagiu T. Sensory polymyeloradiculopathy associated with Toscana virus infection. *J Neurovirol.* 2013;19:508–10. [PubMed](#) <https://doi.org/10.1007/s13365-013-0201-y>
9. Anagnostou V, Papa A. Seroprevalence of Toscana virus among residents of Aegean Sea islands, Greece. *Travel Med Infect Dis.* 2013;11:98–102. [PubMed](#)  
<https://doi.org/10.1016/j.tmaid.2012.11.006>
10. Dupouey J, Bichaud L, Ninove L, Zandotti C, Thirion-Perrier L, de Lamballerie X, et al. Toscana virus infections: a case series from France. *J Infect.* 2014;68:290–5. [PubMed](#)  
<https://doi.org/10.1016/j.jinf.2013.11.006>
11. Magurano F, Baggieri M, Gattuso G, Fortuna C, Remoli ME, Vaccari G, et al. Toscana virus genome stability: data from a meningoencephalitis case in Mantua, Italy. *Vector Borne Zoonotic Dis.* 2014;14:866–9. [PubMed](#) <https://doi.org/10.1089/vbz.2014.1668>
12. Charrel RN. Toscana virus infection. In: Ergonul O, Can F, Akova F, Madoff L, editors. *Emerging infectious diseases*. New York: Academic Press; 2014. p. 111–119.



13. Sanchez P, Vázquez González A. Emergence of novel viruses (Toscana, Usutu) in population and climate change. In: Singh SK, editor. Viral infections and global change. New York: John Wiley & Sons; 2014. p. 535–5.
14. Pietrantoni A, Fortuna C, Remoli ME, Ciufolini MG, Superti F. Bovine lactoferrin inhibits Toscana virus infection by binding to heparan sulphate. *Viruses*. 2015;7:480–95. [PubMed](#)  
<https://doi.org/10.3390/v7020480>
15. Howell BA, Azar MM, Landry ML, Shaw AC. Toscana virus encephalitis in a traveler returning to the United States. *J Clin Microbiol*. 2015;53:1445–7. [PubMed](#)  
<https://doi.org/10.1128/JCM.03498-14>
16. Osborne JC, Khatamzas E, Misbahuddin A, Hart R, Sivaramakrishnan A, Breen DP. Toscana virus encephalitis following a holiday in Sicily. *Pract Neurol*. 2016;16:139–41. [PubMed](#)  
<https://doi.org/10.1136/practneurol-2015-001265>
17. Piacenti S, La Ferla L, Capocardo B, Tosto S, Grasso RS, D'Agata V, et al. A case of meningitis caused by Toscana virus. *Infect Dis Trop Med*. 2016;2:e314.
18. Temocin F, Sari T, Tulek N. Sandfly fever with skin lesions: a case series from Turkey. *J Arthropod Borne Dis*. 2016;10:608–12. [PubMed](#)
19. Veater J, Mehedi F, Cheung CK, Nabarro L, Osborne J, Wong N, et al. Toscana virus meningo-encephalitis: an important differential diagnosis for elderly travellers returning from Mediterranean countries. *BMC Geriatr*. 2017;17:193. [PubMed](#) <https://doi.org/10.1186/s12877-017-0593-2>
20. Pierro A, Ficarelli S, Ayhan N, Morini S, Raumer L, Bartoletti M, et al. Characterization of antibody response in neuroinvasive infection caused by Toscana virus. *Clin Microbiol Infect*. 2017;23:868–73. [PubMed](#) <https://doi.org/10.1016/j.cmi.2017.03.017>
21. Tschumi F, Schmutz S, Kufner V, Heider M, Pigny F, Schreiner B, et al. Meningitis and epididymitis caused by Toscana virus infection imported to Switzerland diagnosed by metagenomic sequencing: a case report. *BMC Infect Dis*. 2019;19:591. [PubMed](#)  
<https://doi.org/10.1186/s12879-019-4231-9>
22. Ayhan N, Charrel RN. An update on Toscana virus distribution, genetics, medical and diagnostic aspects. *Clin Microbiol Infect*. 2020;26:1017–23. [PubMed](#)  
<https://doi.org/10.1016/j.cmi.2019.12.015>

23. Arden KE, Heney C, Shaban B, Nimmo GR, Nissen MD, Sloots TP, et al. Detection of Toscana virus from an adult traveler returning to Australia with encephalitis. *J Med Virol*. 2017;89:1861–4. [PubMed https://doi.org/10.1002/jmv.24839](https://doi.org/10.1002/jmv.24839)
24. Calisher CH, Weinberg AN, Muth DJ, Lazouck JS. Toscana virus infection in United States citizen returning from Italy. *Lancet*. 1987;1:165–6. [PubMed https://doi.org/10.1016/S0140-6736\(87\)92005-8](https://doi.org/10.1016/S0140-6736(87)92005-8)
25. Cordey S, Bel M, Petty TJ, Docquier M, Sacco L, Turin L, et al. Toscana virus meningitis case in Switzerland: an example of the ezVIR bioinformatics pipeline utility for the identification of emerging viruses. *Clin Microbiol Infect*. 2015;21:387.e1–4. [PubMed https://doi.org/10.1016/j.cmi.2014.11.010](https://doi.org/10.1016/j.cmi.2014.11.010)
26. Defuentes G, Rapp C, Imbert P, Durand J-P, Debord T. Acute meningitis owing to *phlebotomus* fever Toscana virus imported to France. *J Travel Med*. 2005;12:295–6. [PubMed https://doi.org/10.2310/7060.2005.12512](https://doi.org/10.2310/7060.2005.12512)
27. Dobler G, Treib J, Haass A, Frösner G, Woesner R, Schimrigk K. Toscana virus infection in German travellers returning from the Mediterranean. *Infection*. 1997;25:325. [PubMed https://doi.org/10.1007/BF01720413](https://doi.org/10.1007/BF01720413)
28. Dominati A, Sap L, Vora S. [Fever in a returning traveler from Tuscany]. *Rev Med Suisse*. 2018;14:294–6. [PubMed https://doi.org/10.1007/BF01720413](https://doi.org/10.1007/BF01720413)
29. Ehrnst A, Peters CJ, Niklasson B, Svedmyr A, Holmgren B. Neurovirulent Toscana virus (a sandfly fever virus) in Swedish man after visit to Portugal. *Lancet*. 1985;1:1212–3. [PubMed https://doi.org/10.1016/S0140-6736\(85\)92886-7](https://doi.org/10.1016/S0140-6736(85)92886-7)
30. Epelboin L, Hausfater P, Schuffenecker I, Riou B, Zeller H, Bricaire F, et al. Meningoencephalitis due to Toscana virus in a French traveler returning from central Italy. *J Travel Med*. 2008;15:361–3. [PubMed https://doi.org/10.1111/j.1708-8305.2008.00221.x](https://doi.org/10.1111/j.1708-8305.2008.00221.x)
31. Gabriel M, Resch C, Günther S, Schmidt-Chanasit J. Toscana virus infection imported from Elba into Switzerland. *Emerg Infect Dis*. 2010;16:1034–6. [PubMed https://doi.org/10.3201/eid1606.091763](https://doi.org/10.3201/eid1606.091763)
32. Howell BA, Azar MM, Landry ML, Shaw AC. Toscana virus encephalitis in a traveler returning to the United States. *J Clin Microbiol*. 2015;53:1445–7. [PubMed https://doi.org/10.1128/JCM.03498-14](https://doi.org/10.1128/JCM.03498-14)

33. Imirzalioglu C, Schaller M, Bretzel RG. [Sandfly fever Naples virus (serotype Toscana) infection with meningeal involvement after a vacation in Italy]. *Dtsch Med Wochenschr.* 2006;131:2838–40. [PubMed](#) <https://doi.org/10.1055/s-2006-957210>
34. Jaijakul S, Arias CA, Hossain M, Arduino RC, Wootton SH, Hasbun R. Toscana meningoencephalitis: a comparison to other viral central nervous system infections. *J Clin Virol.* 2012;55:204–8. [PubMed](#) <https://doi.org/10.1016/j.jcv.2012.07.007>
35. Karunaratne K, Davies N. Toscana virus meningitis following a holiday in Elba, Italy. *Br J Hosp Med (Lond).* 2018;79:292. [PubMed](#) <https://doi.org/10.12968/hmed.2018.79.5.292>
36. Kay MK, Gibney KB, Riedo FX, Kosoy OL, Lanciotti RS, Lambert AJ. Toscana virus infection in American traveler returning from Sicily, 2009. *Emerg Infect Dis.* 2010;16:1498–500. [PubMed](#) <https://doi.org/10.3201/eid1609.100505>
37. Kuhn J, Bewermeyer H, Hartmann-Klosterkoetter U, Emmerich P, Schilling S, Valassina M. Toscana virus causing severe meningoencephalitis in an elderly traveller. *J Neurol Neurosurg Psychiatry.* 2005;76:1605–6. [PubMed](#) <https://doi.org/10.1136/jnnp.2004.060863>
38. Oechtering J, Petzold GC. Acute hydrocephalus due to impaired CSF resorption in Toscana virus meningoencephalitis. *Neurology.* 2012;79:829–31. [PubMed](#) <https://doi.org/10.1212/WNL.0b013e3182661fla>
39. Osborne JC, Khatamzas E, Misbahuddin A, Hart R, Sivaramakrishnan A, Breen DP. Toscana virus encephalitis following a holiday in Sicily. *Pract Neurol.* 2016;16:139–41. [PubMed](#) <https://doi.org/10.1136/practneurol-2015-001265>
40. Pauli C, Schwarz TF, Meyer CG, Jäger G. [Neurological symptoms after an infection by the sandfly fever virus]. *Dtsch Med Wochenschr.* 1995;120:1468–72. [PubMed](#) <https://doi.org/10.1055/s-2008-1055501>
41. Schultze D, Korte W, Rafeiner P, Niedrig M. First report of sandfly fever virus infection imported from Malta into Switzerland, October 2011. *Euro Surveill.* 2012;17:4–6. [PubMed](#) <https://doi.org/10.2807/es.e17.27.20209-en>
42. Schwarz TF, Gilch S, Jäger G. Travel-related Toscana virus infection. *Lancet.* 1993;342:803–4. [PubMed](#) [https://doi.org/10.1016/0140-6736\(93\)91568-7](https://doi.org/10.1016/0140-6736(93)91568-7)
43. Sonderegger B, Hächler H, Dobler G, Frei M. Imported aseptic meningitis due to Toscana virus acquired on the island of Elba, Italy, August 2008. *Euro Surveill.* 2009;14:19079. [PubMed](#) <https://doi.org/10.2807/es.e14.01.19079-en>

44. Tschumi F, Schmutz S, Kufner V, Heider M, Pigny F, Schreiner B, et al. Meningitis and epididymitis caused by Toscana virus infection imported to Switzerland diagnosed by metagenomic sequencing: a case report. *BMC Infect Dis.* 2019;19:591. [PubMed](#) <https://doi.org/10.1186/s12879-019-4231-9>
45. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009;6:e1000097. [PubMed](#) <https://doi.org/10.1371/journal.pmed.1000097>